

Measurement of longitudinal single spin asymmetry of $W^{\pm} \rightarrow \mu^{\pm}$


at forward/backward rapidities with PHENIX at √s = 500 - 510 GeV polarized p+p collisions

Hideyuki Oide (University of Tokyo / RIKEN)
on behalf of PHENIX collaboration


Introduction

Decomposing the spin of the proton w/ parton picture


$$\frac{1}{2} = \underbrace{\frac{1}{2} \sum_{f} (\Delta q_f + \Delta \bar{q}_f)}_{f} + \underbrace{\Delta g}_{f} + \underbrace{L}_{no \ measurements}_{no \ measurements}$$
 Revealing to be small (RHIC) -25% in total (DIS) ("The spin crisis")



Looking into the detail of quark flavor...



Flavor-separating measurements of sea quark's polarized PDF have been desired.




Probing sea quark via W[±]→ ℓ [±]

Longitudinally polarized p+p at \sqrt{s} = 500 (510) GeV at RHIC

Single Spin Asymmetry of W production via parity violating weak coupling

$$A_L = \frac{d\sigma^{\Rightarrow} - d\sigma^{\Leftarrow}}{d\sigma^{\Rightarrow} + d\sigma^{\Leftarrow}}$$

is a flavor-sensitive probe to sea quark polarization.



$$d\sigma(p^{\Rightarrow}p \to W^+) \propto u_L^{\Rightarrow}(x_1)\bar{d}_R(x_2) + \bar{d}_R^{\Rightarrow}(x_1)u_L(x_2)$$

PH ENIX: suitable for leptonic decays of W

- Central Arm:

$$W^\pm o e^\pm, \ |\eta| < 0.35$$
 (next talk)


- Forward Muon Arms:


$$W^{\pm} \to \mu^{\pm}$$

-2.2 < η < -1.2, 1.2 < η < 2.4


via measuring single lepton's pT and rapidity

⇒ Various rapidity regions covered.







RHIC




RHIC




RHIC

























- BBC trigger: "minimum-bias collision" + timing w/ Cerenkov counters


- MulD trigger: conventional, requires "deep" hit pattern for muons in MulD







- BBC trigger: "minimum-bias collision" + timing w/ Cerenkov counters


- MuID trigger: conventional, requires "deep" hit pattern for muons in MuID







- BBC trigger: "minimum-bias collision" + timing w/ Cerenkov counters
- MulD trigger: conventional, requires "deep" hit pattern for muons in MulD






- BBC trigger: "minimum-bias collision" + timing w/ Cerenkov counters
- MulD trigger: conventional, requires "deep" hit pattern for muons in MulD






- BBC trigger: "minimum-bias collision" + timing w/ Cerenkov counters

- MulD trigger: conventional, requires "deep" hit pattern for muons in MulD







- BBC trigger: "minimum-bias collision" + timing w/ Cerenkov counters

- MuID trigger: conventional, requires "deep" hit pattern for muons in MuID

- Mutrg & RPC: snatch online hit patterns in MuTr/RPC ⇒ compare with sagitta trigger maps







- BBC trigger: "minimum-bias collision" + timing w/ Cerenkov counters

- MuID trigger: conventional, requires "deep" hit pattern for muons in MuID

- Mutrg & RPC: snatch online hit patterns in MuTr/RPC ⇒ compare with sagitta trigger maps







- BBC trigger: "minimum-bias collision" + timing w/ Cerenkov counters

- MuID trigger: conventional, requires "deep" hit pattern for muons in MuID

- Mutrg & RPC: snatch online hit patterns in MuTr/RPC ⇒ compare with sagitta trigger maps







- BBC trigger: "minimum-bias collision" + timing w/ Cerenkov counters


- MuID trigger: conventional, requires "deep" hit pattern for muons in MuID

- Mutrg & RPC: snatch online hit patterns in MuTr/RPC ⇒ compare with sagitta trigger maps







Trigger performances (Mutrg)




Turn-on curve





Plateau efficiency

Forward Muon Analysis



Backgrounds:

- Heavy flavor, onium (true muon, irreducible)
- "Fake high pT" caused by decayed hadrons


Tight cuts are applied for "consistency of true high pT muon".

- small multiple scattering : MuTr/MuID/RPC matching

- vertex requirement : Track/vertex(BBC) matching

- timing: RPC

Forward Muon Analysis



Signal: high pT single muon

missing E_T isolation

Backgrounds:

- Heavy flavor, onium (true muon, irreducible)
- "Fake high pT" caused by decayed hadrons


Tight cuts are applied for "consistency of true high pT muon".

- small multiple scattering : MuTr/MuID/RPC matching

- vertex requirement : Track/vertex(BBC) matching

RPC - timing:

Forward Muon Analysis



Backgrounds:

- Heavy flavor, onium (true muon, irreducible)
- "Fake high pT" caused by decayed hadrons

Tight cuts are applied for "consistency of true high pT muon".


- small multiple scattering : MuTr/MuID/RPC matching

- vertex requirement : Track/vertex(BBC) matching

- timing: RPC


Momentum resolution (smearing)

Momentum resolution for cosmic muons




Confirmed the validity of smearing in the simulation with cosmic muons which punch through South&North Arms.

Validation of Cut variables




- Validation of cut positions with J/psi


Cuts in simulation



Single Muon Spectrum

Single muon candidates cross section at tightest cut level





Corrections:

- "SG1xMuIDxBBC" trigger efficiency correction
- Reconstruction x Acceptance correction
- RPC efficiency correction
- Luminosity correction

S/BG was extracted with assuming the W/Z cross section predicted by RHICBOS NLO calculation.


S/BG for muons in 18 - 60 GeV/c range with optimized statistical FOM:


S/BG	North	South		
$\mu+$	0.21	0.40		
μ-	0.42	0.33		

Factor [x0.5 - x2.0] range, as a conservative uncertainty of the S/BG

First W→mu single spin asymmetry (preliminary)

Preliminary approved in March 2012.





- \sqrt{s} = 500 GeV
- Luminosity: ~25 pb-1
- Pol.: ~50%

First forward W asymmetry result

Status quo, and near future


- Finalization of the analysis of the first $W\rightarrow\mu$ data in 2011 is ongoing.
- Took about ~50 pb⁻¹ in 2012 with improved trigger with RPC.
 - Inclusion of VTX/FVTX detectors: better understanding / rejection of BG expected.
- Big data (~250 pb⁻¹) will come up in 2013 with full detector ready. Now proposed for the beam use.


Ye	ear	2011	2012	2013	
Luminosity (pb ⁻¹)	vtx <30cm	~16	30	250?	
	Full vtx	25	49	350?	

Summary

- The first preliminary result of forward arm W→µ single spin asymmetry
- In 2012 run $\sqrt{s} = 510$ GeV run ended successfully.
 - Operation of new muon trigger with RPC3.
 - Taking data with VTX, FVTX, and RPC1.
- In 2013, an integrated luminosity of 250 pb⁻¹ in |vtx| < 30 cm is anticipated with full upgraded hardware set ready.</p>

Expected single spin asymmetry in W→µ at 300 pb⁻¹







Summary

- The first preliminary result of forward arm W→µ single spin asymmetry
- In 2012 run $\sqrt{s} = 510$ GeV run ended successfully.
 - Operation of new muon trigger with RPC3.
 - Taking data with VTX, FVTX, and RPC1.
- In 2013, an integrated luminosity of 250 pb⁻¹ in |vtx| < 30 cm is anticipated with full upgraded hardware set ready.</p>

Expected single spin asymmetry in W→µ at 300 pb⁻¹









Backup

Run9 W→e Result in Central Arm

Run9 \sqrt{s} = 500 GeV integrated luminosity: 8.6 pb⁻¹ in |vtxz| < 30 cm (ERT 4x4b trigger)






Single Spin Asymmetry Result (p_T > 30 GeV/c)

$$W^{+} + Z^{0} : -0.86 \ [-1, -0.56]$$

 $W^{-} + Z^{0} : +0.88 \ [+0.17, +1]$
 $([\] : 68\% C.L.)$

Forward Muon Trigger Upgrade



Motivation of PHENIX forward upgrade project:


Due to limited trigger bandwidth, the original muon trigger "MuID" is short to acquire all W collisions without pre-scaling. ==> Needs a high momentum trigger.

[Strategy]

- Discriminates muon momentum using sagitta
 (Mutrg: parasitic trigger electronics inside MuTr + RPC)
 ==> Trigger p_T > 4 GeV/c
- 2. For ID of spin pattern / bunch crossing at high collision rate, beam beam counter (BBC) is not tolerable. ==> Introduce resistive plate chambers (RPC)
- **3.** Introduce additional hadron absorbers to reject hadronic backgrounds.
- 4. FVTX detector will work for increasing analysis power.

Events/(1 GeV/c)

PHENIX Forward Upgrade Program




Muon Arm Upgrade History

Voor		2009 2010		10	2011		2012			
Year		200)9	2010		2011		2012		
Run Status		pp Run	-	-		pp Run	-	pp Run	-	
FWD	MuT	rg	install/commissioning			operation / Physics Trigger				
	RPC3	S				install	operation Operation Physics Trigg		tion /	
		N		install	commis	sioning			Physics	Physics Trigger
	RPC	1					install	install comm./operat		
	FVT	X						install	comm./operation	
CNIT	HBI)	operation			removed				
CNT	VT	K			install				operation	
W->mu TRG		(MuIDxBBC prescaled)		SG1xMuIDxBBC		SG1xRPC3xBBC				
Luminosity vtx < 30 cm		8.0	6	-		16		30		
√s (GeV)		50	0	_		500		510		

Details of upgrades are spoken on Sep 20, S1-V 16:20 by F. Giordano


Run12 Muon Arm performances

Run12 $\sqrt{s} = 510$ GeV luminosity history




Run12 W→µ trigger performance SG1+RPC3 South


Basic cut w/ DCAr = 3 & RpcDCA = 10



Trigger Rejection Power



Prospects of RPC1 for Run13 Muon Trigger



We can expect factor 2~3 rejection power increase with including RPC1 to the muon trigger in Run13+.

Operational of FVTX detector in Run12

